

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/272151504>

New record and distribution extension of *Leptodactylus paraensis* Heyer, 2005 (Anura, Leptodactylidae) in state of Mato Grosso, Brazil

Article in *Herpetology Notes* · August 2012

CITATIONS

0

READS

217

1 author:



Everton José Almeida

Instituto Federal de Educação, Ciência e Tecnologia de Mato Grosso (IFMT)

11 PUBLICATIONS 99 CITATIONS

SEE PROFILE

New record and distribution extension of *Leptodactylus paraensis* Heyer, 2005 (Anura, Leptodactylidae) in state of Mato Grosso, Brazil.

Ana Bárbara Barros^{1,*}, Domingos de Jesus Rodrigues^{1,2}, Janaina da Costa de Noronha^{1,2}, Everton José Almeida^{1,2}

The large size frog *Leptodactylus paraensis* (Heyer 2005) belongs to *Leptodactylus pentadactylus* group that comprises 19 species (Frost, 2011). *Leptodactylus paraensis* is abundant in native Amazonian forests (Heyer, 2005) and it has been found in the municipality of Claudia, state of Mato Grosso (11°35'9.61" S, 55° 16' 10.86" W, 11° 24' 44.21" S, 55° 19' 25.20" W, 11° 38' 20.40" S, 55° 5' 25.43" W) (Rodrigues et al. 2010), Serra de Kukoinhokren (07°46'S, 51°57'W), state of Pará (Heyer, 2005), and other records from Tapajós National Forest in the western state of Pará also exist, although no voucher specimen was collected (Mendes-Pinto, 2011). Finally, one male of *L. paraensis* was observed during the Rapid Ecological Assessment of Floresta Nacional do Trairão (4° 34' 33.67" S, 55° 24' 14.67" W), in the southwestern portion of the state of Pará (Mendes-Pinto, 2011).

In December 2011 and February 2012 we observed and collected two males of *L. paraensis* (Figure 1) in the southernmost portion of the Amazon rainforest. The specimens were collected in the São Nicolau Farm (09°52'24"S, 58°13'17"W) municipality of Cotriguaçu and in the Parque Estadual do Cristalino (Cristalino State Park) (9° 32' 47" S, 55° 47' 38" W), municipality of Mundo Novo. The São Nicolau Farm is about 10,000 ha, of which 7,500 are of rainforest and are currently protected, while the remaining 2,500 have been reforested (Veloso et al. 1991, Embrapa, 1999). The Cristalino State Park is located at south of Serra do Cachimbo, and the lower Cristalino River is partially bordered from the north to the south by open habitats and it is characterized by a mountainous area with rocky

outcrops protruding amidst the rainforest (Sasaki et al. 2008; Caldwell 2009).

The voucher specimens from São Nicolau Farm and Parque Estadual do Cristalino (Cristalino State Park) have been deposited in the Herpetological Collection of the Federal University of Mato Grosso, Campus of Sinop (ABAM-H 64: Snout-Vent Length (SVL)= 139.33 mm; Head Length (HL)= 62.06mm; ABAM-H 1322: SVL: 128.29mm; HL= 58.42mm). These new records of *L. paraensis* in the state of Mato Grosso extend the species distribution of 1030 Km (ABAM-H 64) and 780 Km (ABAM-H 1322) southwest of the type-locality (Figure 2) in an area threatened by deforestation (Rodrigues et al. 2010). *Leptodactylus paraensis* is not classified as



Figure 1. A. *Leptodactylus paraensis* from São Nicolau Farm (Adult, ABAM-H 64); **B.** *Leptodactylus paraensis* from Parque Estadual do Cristalino (Cristalino State Park) (Adult, ABAM-H 1322).

1 Universidade Federal do Mato Grosso, Acervo Biológico da Amazônia Meridional, Avenida Alexandre Ferronato, 1200, Setor Industrial, CEP 78577-267, Sinop, MT, Brazil

2 Instituto Nacional de Ciências e Tecnologia de Estudos Integrados da Biodiversidade Amazônica – INCT-CENBAM/ CNPq/MCT.

* Corresponding author. e-mail: barros.anabarbara@gmail.com




Figure 2. Schematic map showing the occurrence of *Leptodactylus paraensis* with old records (black filled circles) and the new localities (black filled triangles). The type-locality is represented with a black star.

an endangered species under the list of IUCN Red List of Threatened Species (IUCN, 2012). However, the knowledge on its natural history and distribution is still scarce (Azeredo-Ramos and Gallatti, 2002).

Acknowledgements. We are grateful to Alexandre Farias e Marcelo Lima for helping during the fieldwork, as well as Universidade Federal do Mato Grosso (UFMT), Office National des Forêts (ONF-Brazil) and Secretaria do Meio Ambiente of the state of Mato Grosso (SEMA –MT) for the logistics support. Our thanks goes to Dayana Schmidt Moore for providing comments on an earlier version of this manuscript. Thanks to Fundação de Amparo à Pesquisa do estado do Mato Grosso (FAPEMAT) (proc. n°. 300729/2010) and Ministério da Ciência e Tecnologia / Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (proc. n° 569382/2008-4, 501408/2009-6, 558225/2009-8) for financial support. Fellowship from CNPq to ABB and DJR. We would also like to thank the Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis – (IBAMA) for allowing collection of specimens (n° 10174-1). This is publication 19 in the NEBAM technical series.

References

- Azeredo-Ramos, C., Gallatti, U. (2002): Patterns of amphibian diversity in Brazilian Amazonia: conservation implications. *Biol. Cons.* **103**: 103-111.
- Caldwell A. J. (2009): Amphibians and Reptiles of the Lower Cristalino River Region of the Southern Amazon. Eletronic database. Accessible at: <http://www.snomnh.ou.edu/personnel/herpetology/vitt/Cerrado/Cristalino/index.htm>. Accessed on: Feb 2012.
- Embrapa – Empresa Brasileira de Pesquisa Agropecuária. (1999): Sistema brasileiro de classificação de solos. 1 ed. Rio de Janeiro. Centro Nacional de Pesquisas de Solos. 306 p.
- Frost, D.R. (2011): Amphibian Species of the Word: an online reference. Version 5.4. Accessible at: <http://research.amnh.org/herpetology/amphibia/index.php>. American Museum of Natural History, New York. Accessed on: Feb 2012.
- IUCN, International Union for Conservation of Nature. (2012): Red List of Threatened Species. Version 2011. Accessible at: <http://www.iucnredlist.org>. Accessed on: Feb 2012.
- Heyer, W.R. (2005): Variation and Taxonomic clarification of the large species of the *Leptodactylus* species group (Amphibia, Leptodactylidae) from Central America, Northern South America and Amazonia. *Arquivos de Zoologia*. **3**: 269-348.
- Mendes-Pinto, T.J. (2011): New record and distribution extension of *Leptodactylus paraensis* Heyer, 2005 (Amphibia, Anura, Leptodactylidae) in the state of Pará, Brazil. *Revista de Ciências Ambientais*. **5**: 89-92.
- Rodrigues, D.J., Lima, M.M., Velásquez, C., Mott, T. (2010): Amphibia, Anura, Leptodactylidae, *Leptodactylus paraensis* Heyer, 2005: distribution extension, new state record and geographic distribution map. *Check List*. **3**: 467-46.
- Sasaki, D., Zappi, D., Milliken, W. (2008): Vegetação do Parque Estadual Cristalino (Cristalino State Park), Novo Mundo – MT. Relatório preliminar, programa Flora Cristalino. 53p.
- Veloso, H.P. (1991): Classificação da vegetação brasileira, adaptada a um sistema universal. Rio de Janeiro: IBGE, Departamento de Recursos Naturais e Estudos Ambientais. 123p.

Accepted by Angelica Crottini